

TITLE:

Replicated Dryland Cotton Seeding Rate and Planting Pattern Demonstration, AG-CARES, Lamesa, TX, 2003.

AUTHORS:

Randy Boman, Tommy Doederlein, John Farris, Mark Stelter, and Mark Kelley; Extension Agronomist-Cotton, EA-IPM Dawson/Lynn Counties, CEA-Agriculture Dawson County, Extension Program Specialist-Cotton, Extension Assistant-Cotton.

MATERIALS AND METHODS:

Variety: AFD 3511RR

Experimental design: Randomized complete block with 3 replications

Seeding rate: 2, 4, and 6 seed per row-foot in 40-inch row spacing ( John Deere Max Emerge vacuum planter)

Planting patterns: Each seeding rate was planted in a solid pattern and in a plant 2 rows and skip 1 pattern. For ease of planting, all plots were seeded in a solid pattern and after seedling emergence, cultivator sweeps were used to destroy seedling plants in the skip row.

Plot size: 16 rows by 300 ft long

Planting date: June 11

Weed management: Treflan was applied preplant incorporated at 1.5 pt/acre across all plots on April 30. Roundup WeatherMax herbicide was applied on July 2 at 22 oz/acre with 17 lbs of Ammonia Sulfate. Plots were cultivated on July 14, and August 5.

Rainfall:	April:	0.42"	July:	0.00"
	May:	4.50"	August:	2.29"
	June:	1.80"	September:	1.67"
	Total moisture:	10.68"		

Insecticides: No insecticides were applied at this site. This location is in a active boll weevil eradication zone, but no applications were made by the Texas Boll Weevil Eradication Program.

Fertilizer management: No fertilizers were applied at this site.

Harvest aids: Gramoxone Max was applied at 26 oz/acre on November 17.

Harvest: Plots were harvested on November 19 using a commercial John Deere 7445 with field cleaner bypassed. Harvested material was dumped into a weigh wagon with integral digital scales to determine individual plot weights. Plot yields were adjusted to lb/acre.

Gin turnout:	Grab samples were taken by plot and ginned at the Texas A&M Center at Lubbock to determine gin turnouts.
Fiber analysis:	Lint samples were submitted to the International Textile Center (ITC) at Texas Tech University for HVI analysis, and USDA loan values were determined for each variety by plot.
Ginning costs and seed values:	Ginning costs were based on \$2.25 per cwt. of bur cotton and \$125/ton for seed value. Ginning costs do not include checkoff.
Seed costs:	Seed costs were based on the 2, 4, and 6 seed per row-foot and the 2 x 1 skip row pattern (66.6% of solid planting rate).

## RESULTS AND DISCUSSION:

Significant differences were not observed for some characteristics measured (Tables 1 and 2). Lint turnout ranged from 25.1% to 26.6%. Lint yields varied from a low of 345 lb/acre (6 seed/row-ft solid planting) to a high of 389 lb/acre (4 seed/row-ft solid planting). Generally speaking, lint yields tended to decrease with higher seeding rates regardless of planting pattern. Lint loan values varied from a low of \$0.4805/lb (6 seed/row-ft solid planting) to a high of \$0.5360/lb (2 seed/row-ft 2x1 planting pattern). Higher seeding rates tended to reduce loan value of the harvested lint. Micronaire ranged from a low of 4.8 units (2 seed/row-ft solid and 2x1 planting pattern) to a high of 5.0 units (6 seed/row-ft solid and 2x1 planting pattern). With higher seeding rates, micronaire values were increased. Lower seeding rates produced lower micronaire lint which was not discounted in the loan chart. Staple ranged from 32.8 32nds inch with the 6 seed/row-ft solid to a high of 34.4 with the 4 seed/row-ft 2x1 pattern. Staple was reduced by higher seeding rates due to increased plant competition for water and nutrients. No differences were noted for other fiber properties. After adding lint and seed value, total value/acre ranged from a low of \$203.78 (6 seed/row-ft solid planting) to a high of \$242.85 (2 seed/row-ft solid planting). When subtracting ginning and seed and technology fee costs (Table 3), the net value/acre ranged from a high of \$202.20 (2 seed/row-ft solid planting) to \$149.00 (6 seed/row-ft solid planting), a difference of \$53.20. These data indicate that significant differences were not observed in terms of net value/acre due to number of seed planted per acre. Planting patterns tended to have no effect on overall profitability. However, the 6 seed/row-ft solid planting pattern resulted in excessive competition and reduced yield and fiber quality compared to other seeding rates. It should be noted that no thinning of stands was encountered. Due to excellent weather conditions after planting, and no heavy rainfall and/or high wind events, no substantial stand losses were encountered. In “real world conditions” producers might expect thinning of stands due to sand fighting, environmental damage, etc. Additional multi-site and multi-year applied research is needed to evaluate seeding rates and planting patterns across a series of environments.

## ACKNOWLEDGMENTS:

Appreciation is expressed to Danny Carmichael, Research Associate - AG-CARES; and John Everitt, Research Associate, Texas Agricultural Experiment Station, for their assistance on this project and to Dr. John Gannaway, TAES, Lubbock for his cooperation.

**DISCLAIMER CLAUSE:**

Trade names of commercial products used in this report are included only for better understanding and clarity. Reference to commercial products or trade names is made with the understanding that no discrimination is intended and no endorsement by the Texas A&M University System is implied. Readers should realize that results from one experiment do not represent conclusive evidence that the same response would occur where conditions vary.

Table 1. Results from the replicated dryland cotton seeding rate and planting pattern demonstration, AG-CARES, 2003.

Variety	Lint turnout %	Seed turnout %	Bur cotton yield lb/acre	Lint yield lb/acre	Seed yield lb/acre	Lint loan value \$/lb	Lint value \$/acre	Seed value \$/acre	Total value \$/acre	Ginning cost \$/acre	Seed and tech fee \$/acre	Net value \$/acre
2 seed/ft solid	26.6	46.0	1431	381	659	0.5291	201.65	41.20	242.85	32.21	8.45	202.20 a
2 seed/ft 2x1	25.1	46.2	1459	367	675	0.5360	196.97	42.20	239.17	32.84	5.63	200.70 a
4 seed/ft 2x1	25.4	45.9	1486	378	682	0.5215	197.07	42.67	239.74	33.44	11.26	195.04 a
4 seed/ft solid	25.8	45.3	1503	389	682	0.4998	193.75	42.63	236.38	33.82	16.89	185.66 a
6 seed 2x1	26.1	46.6	1476	386	689	0.4953	191.30	43.06	234.36	33.23	16.89	184.25 a
6 seed solid	26.4	46.3	1306	345	605	0.4805	165.96	37.82	203.78	29.39	25.34	149.00 b
Test average	25.9	46.1	1444	374	665	0.5104	191.12	41.60	232.71	32.49	14.08	186.14
CV, %	6.2	3.4	7.1	7.0	7.0	4.4	7.0	7.1	6.8	7.1	--	7.5
OSL	0.8650	0.9454	0.2831	0.4106	0.3321	0.0859	0.0851	0.3324	0.1056	0.2843	--	0.0087
LSD 0.10	NS	NS	NS	NS	NS	0.034	19.98	NS	23.63	NS	--	20.76

For net value/acre, means within a column with the same letter are not significantly different at the 0.05 probability level.

CV - coefficient of variation, LSD - least significant difference.

OSL - observed significance level, or probability of a great F value.

Note: some columns may not add up due to rounding error.

Assumes:

\$2.25/cwt ginning cost.

\$125/ton for seed.

Value for lint based on CCC loan value from grab samples and ITC HVI results.

Table 2. HVI fiber property results from the replicated dryland cotton seeding rate and planting pattern demonstration, AG-CARES, 2003.

Variety	Micronaire	Staple	Uniformity	Strength	Elongation	Leaf	Rd	+b	Color grade	
	units	32nds inches	%	g/tex	%	grade	reflectance	yellowness	color 1	color 2
2 seed/ft 2x1	4.8	34.1	82.6	30.8	5.4	1.3	75.4	7.5	3.6	1.0
2 seed/ft solid	4.8	34.0	82.4	31.8	5.4	2.3	73.8	7.4	4.0	1.0
4 seed/ft 2x1	4.9	34.4	82.4	31.7	5.4	1.6	74.6	7.3	4.0	1.0
4 seed/ft solid	4.9	33.3	81.9	31.6	5.4	2.0	75.1	7.5	4.0	1.0
6 seed 2x1	5.0	33.6	82.3	30.4	5.6	2.0	74.5	7.4	4.0	1.0
6 seed solid	5.0	32.8	81.1	30.3	5.7	1.6	74.7	7.3	4.0	1.0
Test average	4.9	33.7	82.1	31.1	5.5	1.8	74.7	7.4	3.9	1.0
CV, %	2.1	1.4	0.5	2.4	4.3	31.4	1.1	3.3	5.9	--
OSL	0.1428	0.0232	0.0098	0.1137	0.5166	0.4180	0.3747	0.8686	0.4651	--
LSD 0.10	NS	0.7	0.6	NS	NS	NS	NS	NS	NS	--

CV - coefficient of variation.

OSL - observed significance level, or probability of a greater F value.

LSD - least significant difference.

Table 3. Seed costs for the replicated dryland cotton seeding rate and planting pattern demonstration, AG-CARES, 2003.

AFD 3511R 4450 seed/lb	Number of seed/50 lb bag	Total seed/acre	Number of acres/bag	Seed and tech fee \$/bag	Seed and tech fee \$/field acre
2 seed/ft solid	222,500	26136	8.51	71.90	8.45
4 seed/ft solid	222,500	52272	4.26	71.90	16.89
6 seed/ft solid	222,500	78408	2.84	71.90	25.34
2 seed/ft 2x1 skip	222,500	17422	12.77	71.90	5.63
4 seed/ft 2x1 skip	222,500	34845	6.39	71.90	11.26
6 seed/ft 2x1 skip	222,500	52267	4.26	71.90	16.89
13068 row-ft/acre for 40" rows					seed drop on 2x1 skip uses a 0.6666 factor