



2013 Sorghum Pointers for use of Huskie Herbicide

Compiled by Calvin Trostle, Ph.D.

Texas A&M AgriLife Extension Service—Lubbock

ctrostle@ag.tamu.edu, 806-746-6101

February 2013

A photograph of a lush green field of grain sorghum. The plants are densely packed and have long, upright stalks with small, developing seed heads. The background shows a line of trees under a clear sky. Overlaid on the image is yellow text.

**Weed Management in Grain
Sorghum—New in 2012**

**Huskie Herbicide
Bayer Crop Science**

Sources of Label Information

& **AgriLife** Extension Weed Scientists

- Labels for herbicides, insecticides, fungicides, seed treatments, growth regulators, etc.—access through <http://www.cdms.net>, click 'Services' then 'Labels' then enter Brand name
 - After 'Labels' you can also search by active ingredient (looking for a generic?) through "Other Search Options" but will need to register for a free password
- **Texas High Plains**—Dr. Pete Dotray, Lubbock, (806) 746-6101, pdotray@ag.tamu.edu
- **Central & South Texas**—Dr. Paul Baumann, College Station, (979) 845-3041, pbaumann@ag.tamu.edu
- **North Texas**—Dr. Curtis Jones, Commerce, (903) 886-5354, curtis_jones@tamu-commerce.edu

Huskie Herbicide for GS (2013)



- Selective Post-emerge, including Palmer amaranth, redroot pigweed, kochia, species of morningglory, devil's claw, henbit, mareetail; partial control on bindweed, puncturevine
- Best weed control from label suggests spray by 4" tall weeds

Huskie in Cotton Production Regions



- Huskie appears to be a much better all around option than 2,4-D or dicamba in regions where cotton is grown, particularly where either chemical can severely injure cotton due to drift
 - Huskie does not volatilize, but must physically drift
 - 2,4-D amine is better for minimizing drift, but may give up to 10% less weed control than ester formulations (Brent Bean, TX AgriLife, Amarillo)
- Huskie cost relative to 2,4-D or dicamba may be a concern for some dryland farmers

Huskie Herbicide for GS (2013)

- Apply over-the-top, 3-leaf stage to 12" tall (likely 7-8 leaf stage, or ~4 weeks after planting)
- Pyrasulfotole + 2 active ingredients similar or same as Buctril (bromoxynil)
 - Buctril is already labeled in grain sorghum
- Many tank mix options, but for grain sorghum the key is **atrazine: 0.25-1.0 lbs. (0.5-2.0 pints) atrazine per acre** to 'strengthen and expand weed control' (from the label)
 - This atrazine is somewhat lower than for straight atrazine applications (~1/4 to 1/2 less)

Atrazine with Huskie #1

Especially in Sandy Soils



- Sandier soils especially Lubbock and south require reduced atrazine rates (1.0 to 1.5 pints/A), thus the strong suggestion to include atrazine with Huskie for improved control could elevate atrazine rates too much for sandier soils or raise concerns about rotation to cotton
- Some producers switch to propazine for Pre-plant or Pre-emerge (not labeled POST) due to coarse soil or potential carryover concerns (specifically cotton)

Atrazine with Huskie #2



- Should atrazine applied either pre-plant/pre-emerge be delayed to instead couple with Huskie?
- Probably not—It is always good to do what we can to minimize weeds from establishing in the first place (using PP, PRE), so maintaining early atrazine applications, if that has been your practice is advised (or see other options below, which may include mixes with reduced PP/PRE atrazine rates)

Atrazine with Huskie #3



- The time from early application (pre-plant or pre-emerge) to [Huskie + atrazine] POST applied at sorghum ~12" tall could be up to 4 weeks
- PP or PRE atrazine rates might be reduced to 0.5 lb./A (1 pint/A) in order to permit at least 0.25 lb./A (0.5 pint/A) to use with Huskie for synergism in control
- But other PP or PRE options may be more appropriate...


Atrazine with Huskie #4

Early Pre-plant or Pre-emerge options...

- Using other Pre-plant or Pre-emerge options (including propazine) may be better to provide **preventive weed control** and “save” atrazine for pairing with Huskie
- These options include:
 - ❖ Propazine (Milo-Pro) as PP or PRE
 - ❖ Individually, s-metolachlor (Dual Magnum), alachlor (Micro-Tech, Intro), or acetochlor (Warrant)
 - ❖ s-metolachlor or alachlor in combination with reduced rates of atrazine (e.g., Bicep II Magnum, Cinch ATZ, Bullet, Lariat, etc.)
 - ❖ Also dimethenamid (Outlook), possibly mixed with atrazine (Guardsman Max)
 - ❖ Saflufenacil (Sharpen), possibly mixed with dimethenamid or reduced atrazine rates

Current Bayer Suggestion

Texas High Plains; confirm in other Texas regions



- For optimization of control:
 - 1 pint/A (16 oz.) of Huskie
 - + 1 pint/A (16 oz.) of atrazine
 - + 1 lb./A of Ammonium Sulfate
- Growers on loamy and clayey soils with 2 pints/A rates of atrazine can split the atrazine into two 1 pint/A applications.

Huskie Herbicide for GS (2013)

- Other tank mixes for broadleaf control: includes Ally, dicamba, 2,4-D, Peak, Starane
- Label suggests spray grade ammonium sulfate (AMS) at 0.5-1.0 lbs./A
- NIS (non-ionic surfactant) if tank mix partner requires it
- TX AgriLife has noted only minor to ~15% leaf burn/injury at 4-leaf sorghum, essentially all disappears by 3 weeks; similar to none at 8-leaf stage
- No maturity delay or reduced yield due to Huskie has been observed in TX AgriLife trials, 2009-2011

Huskie Control of P. Amaranth

Hale County, Texas

A



B



Credit: Dr. Pete Dotray

(A) Non-treated control, and (B) good Palmer Amaranth control with Huskie + atrazine.

Huskie Herbicide for GS (2013)



- The overarching advantage of Huskie over 2,4-D or dicamba is reduced sorghum plant injury.
- 2,4-D or dicamba might be considered as a tank mix with Huskie in two possible circumstances:
 - Weeds have moved beyond the 4" tall growth stage (harder to control)
 - If you are dealing with triazine or glyphosate resistant weeds you may need to consider a stronger level of control to further minimize escapes and kill existing weeds

Huskie—TX AgriLife Results

Dr. Brent Bean, Bushland, TX (2009-2010)



- 91%+ control 7 & 42 DAT (days after treatment) of 3-4" Palmer amaranth when applied alone at 13-16 oz./A
- 95%+ Palmer amaranth control when applied with 0.5 lb. atrazine at 13 oz./A
- Adding 4 oz./A dicamba did not improve control
- 80% control on 18" tall pigweed (unlabeled for this late application to grain sorghum)

Huskie—TX AgriLife Results

Dr. Pete Dotray, Halfway, Hale Co. (2010-2011)

- 2010: Huskie + atrazine, 97% control of 2-4" Palmer amaranth at 41 DAT; 96% control of 6-8" weeds @ 37 DAT
- Slight sorghum injury for all POST treatments, but $\leq 5\%$ at 37-41 DAT unless 2,4-D included
- Tank mix with 2,4-D or dicamba reduced yield
- 2011: Huskie + atrazine controlled P. amaranth 94% or more at 21 DAT though control decreased with time (hot, hot year).

Huskie in Crop Rotations



- 1 month: Small grains
- 4 months: alfalfa, millet, grain sorghum, soybean
- 9 months: corn, sunflower
- Cotton? 18 months or field bioassay for reduced restriction
 - Texas A&M AgriLife & Bayer staff have not yet observed a problem rotating to cotton after a Huskie application; label considerations in future may reduce to 9 or 10 months

Huskie & Grain Sorghum Injury



- Sourced from Russ Perkins, Bayer CropScience, Lubbock:
 - Full label rate applications across several grain sorghum nurseries in the Texas High Plains in 2009-2010 noted up to 10% of sorghum lines that demonstrated slightly higher leaf phytotoxicity (injury) at 15-20%, but symptoms were largely gone in < 14 days
 - Breeding staff noted no apparent delay in maturity or reduced yield due to Huskie

Additional Huskie Pointers



- Do not apply Lorsban (chlorpyrifos) with Huskie as unacceptable injury may occur
- Baythroid is a acceptable tank mix insecticide partner
- Do not apply Huskie after mesotrione PRE herbicides (e.g. Callisto, Lumax, Lexar)
- Bayer & crop consultant observations suggest that iron chelate labeled for foliar use may reduce injury potential, particularly for high pH soils (1 lb. of iron chelate per 1 pint of Huskie)

Huskie & Sorghum Forages?

- Caution on sorghum/sudan, sorgo or sweet sorghum, sudangrass (forage sorghum not specifically noted)
- Some hybrid sensitivity? Label states 'not recommended' for use in above sorghum types (unless previously tested on a small area), so producer assumes risk of injury.
- Texas A&M AgriLife does not foresee major Huskie injury potential in any sorghum forage type.
- Bayer's Russ Perkins reports these forages have demonstrated higher leaf burn (20-30% in a few cases), but forage grew out of it.
- Note earlier comment on Fe chelate mix with Huskie

Huskie & Sorghum Seed Production?



- Huskie is acceptable for use in seed production
- This is in contrast to the restrictions for seed production using 2,4-D or dicamba, which can risk sharp reduction in seed production and seed viability

Problems with Huskie?



- Complaints and disappointment have been few
 - Most likely on dryland where weeds were hardened off
 - Weeds were already sizable and control was reduced
- “Best herbicide for grain sorghum in 30 years...”
 - Dr. Brent Bean, former Extension agronomist & weed scientist, Texas A&M AgriLife--Amarillo

Huskie & Weed Resistance

- +++ Glyphosate-resistant Palmer amaranth is here in the TX South Plains
 - Huskie will offer cotton producers a much needed means of diversifying chemistries
- - - - Kansas State reports (Jan. 2013) early Palmer amaranth resistance to Huskie (pyrasulfotole component) where mesotrione herbicides have been used in corn many years
 - HPPD inhibitors—mesotriones, pyrasulfotole
 - Resistance will be slower to develop where use is 1 year out of 3 or 4 years
 - Don't "abuse" the product or rely on it solely

A Final Thought

Huskie + Atrazine POST vs. Good PRE Program

- The advent of Huskie for Texas sorghum should not overshadow the fact that a farmer's weed control decision about sorghum pre-plant/pre-emergent weed control is more important than the option Huskie offers.
- **In fact, pre-emerge (or PRE) weed control is the most important weed control decision a farmer will make in grain sorghum production in Texas.**
- Dr. Wayne Keeling, AgriLife Research weed scientist, Lubbock, notes that although Huskie is a good asset, our priority effort should be focused on effective PRE weed control. **Our goal?**—Prevent weeds in the first place, especially during sorghum emergence and early growth. Then post-emergent (POST) weed control (Huskie, dicamba, Ally, Permit, 2,4-D, etc.) can focus on controlling escapes as needed. Furthermore, if PRE weed control is good, producers may be able to delay needed POST control by 1-3 weeks (if the label is extended past 12" tall) to provide a longer window of either direct control of existing weeds or extending residual control further into the growing season.

Future Huskie Label Consideration #1



- Rotation to cotton on the Huskie label remains at 18 months or field bio-assay.
- Texas A&M AgriLife and Bayer staff have not yet observed any significant rotation issues to cotton
- It is anticipated this rotation restriction may eventually be shortened.

Future Huskie Label Consideration #2

- There is consideration of expanding the Huskie label for applications up to flag leaf emergence. There are two potential uses of Huskie in this situation:
 - 1) Later-season rescue treatment: Though this may be appealing, view this only as a last resort and do not rely on late application if it should be made 1-3 weeks sooner. **Waiting until nearly flag leaf emergence means that pigweed and other problem weeds are larger and much harder to kill.** Weed control will more likely be incomplete and a new round of pigweed seed may be generated.
 - 2) If Pre-emerge weed control is good with only a few escapes, then a later mid-season application may be needed to clean these up. Good PRE weed control might provide the opportunity to delay mid-season Huskie or (Huskie + atrazine) applications past the 12" tall sorghum, which enables control of later emerged weeds and/or extend the opportunity for preventive weed control with atrazine.