

Response of cotton varieties to Bacterial blight (Race 18)



Dr. Terry A. Wheeler
Research Plant Pathologist, Texas A&M AgriLife Research

Dr. Jason E. Woodward, Extension Plant Pathologist
Texas A&M AgriLife Extension Service

TEXAS A&M
AGRI LIFE
RESEARCH

TEXAS A&M
AGRI LIFE
EXTENSION

The information given herein is for educational purposes only. Reference to commercial products or trade names is made with the understanding that no discrimination is intended and no endorsement by the Texas A&M AgriLife Extension Service, Texas A&M AgriLife Research Experiment Station and the Texas A&M System is implied.

Brand	Variety	Bacterial blight rating	Brand	Variety	Bacterial blight rating
All-Tex	All-Tex Apex B2RF	S	Fibermax	FM 2322GL	S
All-Tex	All-Tex Arid B2RF	S	Fibermax	FM 2334GLT	R
All-Tex	All-Tex Concho B2XF	Unk	Fibermax	FM 2484B2F	R
All-Tex	All-Tex Dinero B2RF	S	Fibermax	FM 2989GLB2	R
All-Tex	All-Tex Edge B2RF	S	Fibermax	FM 8270GLB2	R
All-Tex	All-Tex Epic RF	S	Fibermax	FM 9180B2F	R
All-Tex	All-Tex Nitro-44B2RF	R	Fibermax	FM 9250GL	R
Croplan Genetics	CG 3035RF	S	NexGen	NG 1511B2RF	S
Croplan Genetics	CG 3156B2RF	S	NexGen	NG 1572RF	R
Croplan Genetics	CG 3787B2RF	R	NexGen	NG 3306B2RF	S
Deltapine	DP 0912B2RF	S	NexGen	NG 3405B2XF	S
Deltapine	DP 104B2RF	S	NexGen	NG 3406B2XF	S
Deltapine	DP 1044B2RF	PS	NexGen	NG 3500XF	Unk
Deltapine	DP 1048B2RF	S	NexGen	NG 3517B2XF	S
Deltapine	DP 1050B2RF	S	NexGen	NG 4111RF	R
Deltapine	DP 1212B2RF	S	NexGen	NG 4545B2XF	R
Deltapine	DP 1219B2RF	S	NexGen	NG 5007B2XF	S
Deltapine	DP 1252B2RF	S	Phytogen	PHY 222WRF	S
Deltapine	DP 1311B2RF	S	Phytogen	PHY 223WRF	R
Deltapine	DP 1321B2RF	S	Phytogen	PHY 243WRF	R
Deltapine	DP1359B2RF	PR	Phytogen	PHY 308WRF	S
Deltapine	DP 1410B2RF	R	Phytogen	PHY 312WRF	PS
Deltapine	DP 1441RF	S	Phytogen	PHY 333WRF	S
Deltapine	DP 1454NRB2RF	S	Phytogen	PHY 339WRF	R
Deltapine	DP 1518B2XF	Unk	Phytogen	PHY 367ERF	S
Deltapine	DP 1522B2XF	S	Phytogen	PHY 375WRF	R
Deltapine	DP 1538B2XF	S	Phytogen	PHY 417WRF	S
Deltapine	DP 1549B2XF	S	Phytogen	PHY 427WRF	S
Deltapine	DP 1553B2XF	S	Phytogen	PHY 444WRF	S
Deltapine	DP 1555B2RF	S	Phytogen	PHY 487WRF	S
Deltapine	DP 1558NRB2RF	S	Phyhtogen	PHY 495W3RF	S
DynaGro	DG 2615B2RF	PR	Phytogen	PHY 499WRF	S
DynaGro	DG 3109B2XF	S	Stoneville	ST 4747GLB2	S
DynaGro	DG 3544B2XF	Unk	Stoneville	ST 4848GLT	S
Fibermax	FM 1320GL	PS	Stoneville	ST 4946GLB2	S
Fibermax	FM 1830GLT	R	Stoneville	ST 4949GLT	S
Fibermax	FM 1900GLT	PR	Stoneville	ST 5032GLT	S
Fibermax	FM 1911GLT	R	Stoneville	ST 5115GLT	R
Fibermax	FM 1944GLB2	S	Stoneville	ST 5289GLT	R
Fibermax	FM 2007GLT	PR	Stoneville	ST 6182GLT	S
Fibermax	FM 2011GT	R	Stoneville	ST 6448GLB2	R

*S = susceptible (> 90% incidence of bacterial blight in inoculated field trials).

PS = partially susceptible (60 – 80% incidence of blight typically in inoculated field trials).

PR = partially resistant (10 to 25% incidence of blight typically in inoculated field trials).

R = Resistant (< 10% incidence of blight in inoculated field trials).

Unk = known, has not been tested yet in inoculated field trials.